


NNRTI (Non-Nucleoside Reverse Transcriptase Inhibitor) Drug Azt and Their Preparation & Determination a Patent Evaluation of Efavirenz

Abstract

Human immunodeficiency virus type-1 (HIV-1) is the causative agent for the transmission and development of the acquired immunodeficiency syndrome (AIDS). Efavirenz (EFV) a potent NNRTI, is a significant component of a very effective (protease sparing) regimen when co administered with AZT and 3TC (De Aquino Ribeiro, et al. [1]). Importance of Efavirenz led to the development of several strategies for its synthesis, and preparation was described in this paper, the world great potential drug Efavirenz still inaccessible to millions of people because of its atom economy and toxic (Montgomery, et al. [2]). Patent evaluation on process patents and well defined summary of patents. During the course of drug development most of the world pharmaceutical companies attempt well in the handling of environment friendly Efavirenz development especially Merck. We have surprisingly found that compound of Efavirenz can be prepared in good yield and high purity by Cyclisation of compound of (S)-2-(2-amino-5-chlorophenyl)-4-cyclopropyl-1,1,1-trifluorobut-3-yn-2-ol with urea. Finally Analytical method for determination Efavirenz has the absorbance maxima at 561 nm for the first order derivative visible-Spectrophotometric method.

Keywords

HIV-1; AIDS; AZT; DTG Benzoxazinones; Efavirenz; Urea; Cyclisation; Crystallisation; UV; Spectrophotometric method; ART; CPA;



Efavirenz; DMP 266;

Efavirenz; L-743725(+)-enantiomer); DMP-266; L-741211(racemate); L-743726; Stocrin; Sustiva (S)-(-)-6-Chloro-4-(cyclopropylethynyl)-4-(trifluoromethyl)-2,4-dihydro-1H-3,1-benzoxazin-2-One.

Open Access

Research Article

Dr. Krishna Sarmapathy*

Head IPL Research Centre, India

*Address for Correspondence

Krishna sarmapathy, Head IPL Research Centre, India


Submission: April 12, 2019

Published: May 30, 2019


Copyright: ©  This work is licensed under Creative Commons Attribution 4.0 License

Introduction

At the moment, India recommends Efavirenz (EFV) based antiretroviral treatment (ART) to treat HIV, which is currently cheaper than DTG-based ART. However, as pharmaceutical licences on DTG are now lifting in India, generic versions of DTG are likely to bring its cost close to that of EFV. Research from the USA and Europe, where DTG is recommended as the first option for treatment, suggests DTG has more tolerable side effects, can enable people to suppress HIV levels quickly and is less likely to lead to drug resistance. As a result, DTG's use in these countries has substantially reduced the number of people who fail first-line treatment and are lost to care or require a switch to second-line ART. Using a mathematical simulation model of how HIV can advance and how it responds to treatment, the study, published by the Journal of International AIDS Society, examined the clinical and economic impact DTG could have on India's HIV response in comparison to EFV. Researchers found using DTG as the first option for HIV treatment would be cost-effective, and potentially cost-saving, for India's public health system. At 48 weeks, modeling suggests 90% of people on DTG would be virally suppressed compared to 82% of those on EFV. Longer term, DTG was projected to increase the proportion of people living with HIV on treatment who were still alive after five years from 76.7% to 83%. Crucially, researchers found DTG had the potential to avert 13,000 new HIV infections over five years, equivalent to a 20% decrease in HIV incidence, resulting in a saving of US\$800,000 in projected care costs. When all factors are taken into account, care costs for the number of people initiating treatment on EFV were estimated to be US \$139 million at two years and US \$590 million at five years. In comparison, the care costs for DTG at two years was found to be lower at US \$137 million and cost-neutral at five years at US \$590 million. Cost increases linked to DTG were identified by the model, although these reflect the fact that more people are likely to live longer when on DTG, incurring greater lifetime care costs.



ketone: Asymmetric synthesis of the reverse transcriptase inhibitor L-743,726. *Tetrahedron Lett* 1995,36,(49):8937-40.



EP 1332757 A1

When a commercial market already exists for the RMs used in synthesizing an API, their cost can be rather modest. When RMs used in synthesizing an API have no other commercial use, however, they can contribute very substantially to API cost. With a continued growth of volume demand, improved chemistry and competition from multiple suppliers, however, the cost of API RMs can greatly decrease over time. The inhibitor of HIV-1 RT, EFV, provides an illustration of this situation. Cyclopropylacetylene (CPA) is an RM for the synthesis of EFV (Figure 4). During clinical trials, when the demand for CPA was only a few metric tons, this material was produced at a price of USD800-1,350/kg. When the drug was first approved in 1998, and demand for CPA was about 50 metric tons per year, the price of CPA had fallen to USD350/kg. Today, with global demand for EFV at greater than 1,000 metric tons/year, CPA can be purchased for about USD50-60/kg. In the earliest stages of production, nearly 1 kg of CPA was needed to produce a kilogram of EFV. Current production processes are more efficient; roughly 3 kg of EFV is now produced for each 1 kg of CPA used. From this it can be roughly estimated that the contribution of CPA to the cost of EFV API production has fallen from as high as USD425/kg to about USD17-20/kg today.

The most recent chemistry for asymmetric alkylation of manufacturing EFV uses inexpensive, safe reagents and processing at ambient temperature to reach EFV pricing that would have been thought impossible when the drug was launched by Dupont Pharmaceuticals in 1998 Biao J, Yugui S. inventors; Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, assignee. Amino alcohol ligand and its use in preparation of chiral propargylic tertiary alcohols and tertiary amines via enantioselective addition reaction. US Patent 7,439,400. 2008 October 21. Bollu RB, Ketavarapu NR, Indukuri VSK, Gorantla SR, Chava S. inventors; Laurus Labs Private Limited, assignee. Efficient process to induce enantioselectivity in procarbonyl compounds. US Patent

April 1997. Efavirenz has been obtained by two related ways: 1) The acylation of 4-chloroaniline (I) with pivaloyl chloride (II) by means of Na₂CO₃ in toluene gives the expected anilide (III), which is acylated with ethyl trifluoroacetate by means of butyllithium in THF yielding, after hydrolysis with HCl, 2'-amino-5'-chloro-2,2,2-trifluoroacetophenone (IV). The benzylation of (IV) with 4-methoxybenzyl chloride (V) in basic alumina affords the protected acetophenone (VI), which is regioselectively condensed with cyclopropylacetylene (VII) [obtained by cyclization of 5-chloro-1-pentyne (VIII) by means of butyllithium in cyclohexane] by means of butyllithium in THF in the presence of (1R,2S)-1-phenyl-2-(1-pyrrolidinyl)-1-propanol (IX) giving the (S)-isomer of the tertiary alcohol (X) exclusively. The cyclization of (X) with phosgene and triethylamine or K₂CO₃ in toluene/THF yields the benzoxazinone (XI), which is finally deprotected with ceric ammonium nitrate in acetonitrile/water. 2) The condensation of 2'-amino-5'-chloro-2,2,2-trifluoroacetophenone (IV) with cyclopropylacetylene (VIII) by means of butyllithium or ethylmagnesium bromide in THF gives (?)-2-(2-amino-5-chlorophenyl)-4-cyclopropyl-1,1,1-trifluoro-3-butyn-2-ol (XII). The cyclization of (XII) with carbonyldiimidazole (XIII) in hot THF yields the racemic benzoxazinone (XIV). Compound (XIV) is submitted to optical resolution by condensation with (S)-(-)-camphanoyl chloride by means of dimethylaminopyridine (DMAP) in dichloromethane to give the acyl derivative (XVI) as a diastereomeric mixture that is resolved by crystallization and finally decomposed with HCl in ethanol or butanol. Corley, E.G.; Thompson, A.S.; Huntington, M.F.; Grabowski, E.J.J.; Use of an ephedrine alkoxide to mediate enantioselective addition of an acetylide to a prochiral

Application 2012/0264933 A1. 2012 October 18. FPPs for adult ART are usually capsules or tablets. A general rule-of-thumb is that an FPP as a conventional, solid oral dosage formulation costs about 33-40% more than the corresponding API in a competitive market. It has been widely quoted, conversely, that APIs contribute about 60-80% of the cost of an FPP. The API contribution to FPP cost increases with the complexity of synthesis and API cost per kilogram. Although marketing is a substantial incremental cost for originator pharmaceutical companies, generic producers do not incur high marketing costs for ART.

Example 1:




1a

Materials	Amount	Mol	MW
Ketone 1a	1.00 kg	4.47	223.58
(1R, 2S)-N-pyrrolidinyl norephedrine	1.35 kg	6.58	205.30
cyclopropyl acetylene	361.9 g	5.47	66.10
n-BuMgCl (2.0 M in THF)	2.68 L	5.37	
2,2,2-trifluoroethanol (99%)	429.5 g	4.29	100.04
ZnEt ₂ (0.892 M in hexane)	6.02 L	5.37	
THF	9.36 L		
30% K ₂ CO ₃	550 mL		
30% citric acid	2.0 L		
Toluene (for crystallization, 2 mL/g of 4)	2.6 L		
Heptane (for crystallization, 4 mL/g of 4)	5.2 L		

To a solution of trifluoroethanol and (1R, 2S)-N-pyrrolidinyl norephedrine in THF (9L) under nitrogen is added a solution of diethylzinc in hexane at 0°C slowly enough to keep the temperature below 30°C. The mixture is stirred at room temperature for 0.5~1 h. In another dry flask a solution of chloromagnesiumcyclopropylacetylide is prepared as follows: To neat cyclopropyl acetylene at 0°C is added a solution of n-butylmagnesium chloride slowly enough to keep the internal temperature <30°C. The solution is stirred at 0°C for ~40 min and transferred to the zinc reagent via cannula with 0.36 L of THF as a wash. The mixture is cooled to -10 °C and ketoaniline 1a is added. The mixture is stirred at -2 to -8 °C for 35 h, warmed to room temperature, stirred for 3 h, and quenched with 30% potassium carbonate over 1.5 h. The mixture is stirred for 4 h and the solid is removed by filtration and washed with THF (2 cake volume). The wet solid still contains ~18 wt% of pyrrolidinyl norephedrine and is saved for further study. The filtrate and wash are combined and treated with 30% citric acid. The two layers are separated. The organic layer is washed with water (1.5 L). The combined aqueous layers are extracted with 2.5 L of toluene and saved for norephedrine recovery. The toluene extract is combined with the organic solution and is concentrated to ~ 2.5 L. Toluene is continuously feeded and

distilled till THF is not detectable by GC. The final volume is controlled at 3.9 L. Heptane (5.2 L) is added over 1 h. The slurry is cooled to 0 °C, aged for 1 h, and filtered. The solid is washed with heptane (2 cake volume) and dried to give 1.234 Kg (95.2% yield) of amino alcohol 3 as a white crystalline. The material is 99.8 A% pure and 99.3% ee.


Example 2:



	FW	g	mL	mmol	equiv
amino alcohol 3	289	100		346	1
4-nitrophenylchloroformate	201.6	73.2		363	1.05
KHCO ₃	100	45		450	1.3
2N KOH	56		346	692	2.0
H ₂ O			654		
MTBE			500		

To a three necked round bottom flask, equipped with a mechanical stirrer, nitrogen line, and thermocouple, was charged the solid amino alcohol 3, MTBE (500 L), and aqueous KHCO₃ (45 g in 654 mL H₂O). Solid 4-nitrophenyl chloroformate was added, in 4 batches, at 25°C. During the addition the solution pH was monitored. The pH was maintained between 8.5 and 4 during the reaction and ended up at 8.0. The mixture was stirred at 20-25°C for two hours. Aqueous KOH (2N) was added over 20 minutes, until the pH of the aqueous layer reached 11.0. The layers were separated and 500 mL brine was added to the MTBE layer. 0.1 N Acetic acid was added until the pH was 6-7. The layers were separated and the organic phase was washed with brine (500 mL). At this point the mixture was solvent switched to EtOH/IPA and crystallized as recited in Examples 5 and 6.


Example 3:



	FW	g	mL	mmol	equiv
amino alcohol 3a	289	100		346	1
phosgene (20 wt% in toluene)	99	41	216	415	1.2
KHCO ₃	100	86.5		865	2.5
H ₂ O			500		
Toluene			500		

To a three necked round bottom flask, equipped with a mechanical stirrer, nitrogen line, and thermocouple, was charged the solid amino alcohol 3a, toluene (500 mL), and aqueous KHCO₃ (86.5 g in 500 mL H₂O). Phosgene solution in toluene was added at 25°C, and the mixture was stirred at 20-25°C for two hours. The layers were separated and the organic phase was washed with brine (500 mL). At this point the mixture was solvent switched to EtOH/IPA and crystallized as recited in Examples 5 and 6.

Example 4:



	FW	g	mL	mmol	equiv
amino alcohol 3a	289	100		346	1
phosgene (gas)	99				
KHCO ₃	100	86.5		865	2.5
H ₂ O			500		
MTBE			500		


To a three necked round bottom flask, equipped with a mechanical stirrer, nitrogen line, and thermocouple, was charged the solid amino alcohol 3a, MTBE (500 mL), and aqueous KHCO₃ (86.5 g in 500 mL H₂O). Phosgene gas was slowly passed into the solution at 25°C, until the reaction was complete. The layers were separated and the organic phase was washed with brine (500 mL). At this point the mixture was solvent switched to EtOH/IPA and crystallized as recited in Examples 5 and 6.

Example 5: Crystallization of efavirenz from 30% 2-Propanol in Water using a ratio of 15 ml solvent per gram efavirenz Using Controlled Anti-Solvent Addition on a 400 g Scale. 400 g. of efavirenz starting material is dissolved in 1.8 L of 2-propanol. The solution is filtered to remove extraneous matter. 1.95 L of deionized (DI) water is added to the solution over 30 to 60 minutes. 10 g. to 20 g. of efavirenz seed (Form II wetcake) is added to the solution. The seed bed is aged for 1 hour. The use of Intermig agitators is preferred to mix the slurry. If required (by the presence of extremely long crystals or a thick slurry), the slurry is wet-milled for 15-60 seconds. 2.25 L of DI water is added to the slurry over 4 to 6 hours. If required (by the presence of extremely long crystals or a thick

slurry), the slurry is wet- milled for 15-60 seconds during the addition. The slurry is aged for 2 to 16 hours until the product concentration in the supernatant remains constant. The slurry is filtered to isolate a crystalline wet cake. The wet cake is washed with 1 to 2 bed volumes of 30% 2-propanol in water and then twice with 1 bed volume of DI water each. The washed wet cake is dried under vacuum at 50°C.

Example 6: Crystallization of efavirenz from 30% 2-Propanol in water using a ratio of 15 ml solvent per gram efavirenz Using a Semi-Continuous Process on a 400 g Scale. 400 g. of efavirenz starting material is dissolved in 1.8 L of 2- propanol. A heel slurry is produced by mixing 20 g. of Form II efavirenz in 0.3 L of 30 % (v/v) 2-propanol in water or retaining part of a slurry from a previous crystallization in the crystallizer. The dissolved batch and 4.2 L of DI water are simultaneously charged to the heel slurry at constant rates over 6 hours to maintain a constant solvent composition in the crystallizer. Use of Intermig agitators during the crystallization is preferred. During this addition the slurry is wet-milled when the crystal lengths become excessively long or the slurry becomes too thick. The slurry is aged for 2 to 16 hours until the product concentration in the supernatant remains constant. The slurry is filtered to isolate a crystalline wet cake. The wet cake is washed with 1 to 2 bed volumes of 30% 2-propanol in water and then twice with 1 bed volume of DI water each. The washed wet cake is dried under vacuum at 50°C.

Example 7: Preparation of Amino Alcohol 3 and eeUpgrading-Through Process,




Materials	Amount	Mol	MW
Ketone 1	1.00 kg	4.47	223.58
(1R, 2S)-N-Pyrrolidinyl norephedrine	1.35 kg	6.58	205.30
Cyclopropyl acetylene	361.9 g	5.47	66.10
n-BuMgCl (2.0 M in THF)	2.68L	5.37	
Trifluoroethanol (99%)	429.5 g	4.29	100.04
ZnEt ₂ (0.892 M in hexane)	6.02L	5.37	
THF	9.36L		
30% H ₂ CO ₃	1.2 L		
1 M Citric acid	3.5 L		
Heptane	12L		
Isopropyl acetate (IPAc)	40L		
12N HCl	405 mL	4.89	
tert-Butyl methyl ether (MTBE)	6L		
Toluene	6.25L		
Na ₂ CO ₃	1.2 kg	11.36	

a solution of diethyl zinc in hexane was added to a solution of trifluoroethanol (429.5g, 4.29'mol) and (1R, 2S)-N-

pyrrolidinylnorephedrine (1.35 kg, 6.58 mol) in THF (9 L), under nitrogen, at 0°C. The resulting mixture was stirred at room temperature for approx 30 min. In another dry flask a solution of chloromagnesium-cyclopropylacetylde was prepared as follows. To a solution of n- butylmagnesium chloride in THF (2 M, 2.68 L, 5.37mol) was added neat cyclopropylacetylene at 0°C keeping the temperature < 25°C. The solution was stirred at 0°C for 1 ~ 2h. The solution of chloromagnesiumcyclopropylacetylde was then warmed to room temperature and was transferred into the zinc reagent via cannula over 5 min followed by vessel rinse with 0.36 L of THF. The resulting mixture was aged at ~ 30°C for 0.5 h and was then cooled to 20°C. The ketoaniline 1 (1.00 kg, 4.47 mol) was added in one portion as a solid, and the resulting mixture was stirred at 20-28°C for 3h. The reaction was quenched with 30% aq. potassium carbonate (1.2 L) and aged for 1 h. The solid waste was filtered and the cake was washed with THF (3 cake volumes). The filtrate and wash were combined and solvent switched to IP Ac. The IPAc solution of product 3 and pyrrolidinylnorephedrine was washed with citric acid (3.5 L) and with water (1.5 L). The combined aqueous layers were extracted with IPAc (2 L) and saved for norephedrine recovery. To the combined organic layers was added 12N HCl (405 mL, 4.88 mol), to form a thin slurry of the amino alcohol-HCl salt. The mixture was aged for 30 min at 25°C and was then dried azeotropically. The slurry was aged at 25°C for 30 min and filtered. The cake was washed with 2.5 L of IPAc and dried at 25°C under vacuum/nitrogen for 24 h to give 1.76 kg of the wet HCl salt. The salt was dissolved in a mixture of MTBE (6 L) and aq Na₂C₂O₃ (1.18 kg in 6.25 L water). The layers were separated and the organic layer was washed with 1.25 L of water. The organic layer was then solvent switched into toluene. Heptane (5 L) was added over 1 h at 25°C. The slurry was cooled to 0°C, aged for 1 h, and filtered. The solid was washed with heptane (2 cake volumes) and was dried to give 1.166 kg (90% overall yield) of amino alcohol 3 as a white crystalline solid.

Norephedrine Recovery

The aqueous solution was basified to pH13 using 50% aq NaOH, and extracted with heptane (2L). The heptane solution was washed with water (1L) and concentrated to remove residual IPAc and water. The final volume was adjusted to about 3L. The heptane solution was cooled to -20°C, aged for 2 h, and filtered. The solid was washed with cold heptane (1 cake volume) and dried to give 1.269 kg solid (94% recovery)



Intermedaite Preparation

The key intermediate of efavirenz: (S)-1-(2-amino-5-chlorophenyl)-1-trifluoromethyl-3-cyclopropyl-2-propyne-1-ol (1) was prepared. p-Chloroaniline. (2) reacted with pivaloyl chloride to give N-(4-chlorophenyl)-2,2-dimethylpropanamide (3). After a Friedel-Crafts acylation of compound 4 catalyzed by aluminumtrichloride, and a hydrolysis under acidic conditions, 4-chloro-2-(trifluoroacetyl) aniline hydrochloride. (4) Was obtained. In this step, the hazardous reagent n-butyllithium was abolished. Compound 5 was basified to afford 4-chloro-2-(trifluoroacetyl) aniline (5). Then compound 6 reacted with chloromagnesiumcyclopropylacetylde under the catalytic system formed by zinc chloride and (1R,2S)-1-phenyl-2-(1-pyrrolidinyl)-1-propanol (6) and a small amount of purified 1 to obtain the target compound 1 by asymmetric autocatalysis with an overall yield of 69.5%(based on 2). In the last step, zinc chloride was used instead of diethylzinc, the yield was increased from 79% to 85.6%. And ligand 7 was recovered with a recovery rate of 98%, which could reduce the cost successfully. This improved synthetic route has mild reaction conditions and the yield of each step was excellent.

Materials and Methods

General procedure-¹H NMR spectra were recorded on a Gemini 300 MHz FT NMR spectrometer; the chemical shifts are reported in δ ppm relative to TMS. The IR spectra were recorded in the solid state as KBr dispersion using Perkin Elmer Spectrophotometer. The mass spectra were recorded on Shimadzu LCMS-QP8000, LC-MS and AB-4000 Q-trap LC-MS/MS.

Representative Procedure for the Preparation of Efavirenz

A solution of (S)-5-chloro-α-(cyclopropylethynyl)-2-amino-α-(trifluoroethyl)benzene methanol (I) (10 gm, 0.034mole) and urea (2.48 gm, 0.0414 mole in toluene are refluxed in the presence of para-toluenesulphonic acid (PTSA) (0.31 gm, 0.0006 mole) for 20 hrs. After completion of the reaction as monitored by Thin Layer Chromatography (TLC), the reaction mixture was cooled to room temperature and then 30 ml of water was added. The obtainedaqueous layer was extracted with toluene. The combined toluene layers were concentrated under reduced pressure. The obtained residue was cooled to room temperature and 30 ml of methanol was added and stirred. Methanol wasdistilled of to remove the toluene traces. Again methanol was added to the residue and cooled to 10-15 °C. To thisresidue water was added at 10-15 °C and stirred for 3 hrs. The obtained white solid was re-crystallized in tolueneand heptane mixture to get 9.2 gms of (4S)-6-chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2H-3,1-benzoxazin-2-one(II) (Efavirenz). ¹H NMR (300MHz in DMCO-d₆): δ 0.782-0.815 (m, 2H), 0.908-0.956 (m,2H), 1.557-1.646 (m, 1H, cyclopropyl), 6.98-7.013(d, 1H), 7.45(s, 1H), 7.55-7.58 (dd, 1H), 11.11(s, 1H).

Mass

Mass m/z; 314.2, in negative ion mode. Yield: 85 %, MR: 136-139°C, SOR -92.8°(c 0.3 in MeOH) Chemicals and reagents Ethanol was used for the quantification of efavirenz which was procured from Sigma Aldrich. EFA film coated tablet of strength 600 mg Sustiva (Bristo-Myers Squibb-Germany) and Efavir (Cipla Ltd-India) were procured from the market. The pure sample of efavirenz for the research work was procured from Shasun Pharmaceuticals Ltd, Pondicherry. Hydrochloric acid, sodium hydroxide, and-naphthol were purchased from Sigma Aldrich. All the chemicals and reagents for the development of new analytical method to estimate efavirenz was of analytical grade. Preparation of -naphthol reagent Accurately 1g of naphthol was weighed and transferred to 100ml volumetric flask. To it 30ml of 0.1N NaOH was added and shaken such that -naphthol gets completely dissolved in NaOH. Make up the volume to 100ml with 0.1N NaOH. Preparation of Efavirenz standard solutions 100 mg of the drug was weighed and transferred in to 100 ml volumetric flask, to it ethanol was added and volume was made up to 100ml with ethanol. From this 10ml was transferred to 100ml volumetric flask and volume was made up to 100ml with ethanol such that the solution gives the concentration of 100 /. Preparation of reagent blank The reagent blank was prepared in the same manner as discussed, omitting the standard drug solution. Procedure for the analysis of tablet formulation Twenty tablets of Efavirenz (600mg) and Sustiva were purchased from local market. Accurately weighed and finely powdered. The weight of the tablet equivalent to 100mg from each brand was accurately weighed and extracted with ethanol. The yellow coloured complex (Scheme-1) having final concentration of 100mg/ml of each brand was prepared. The absorbance of the solutions was measured at 561nm using reagent blank.

Results and Discussion

Selection of analytical wavelength: Appropriate dilutions were prepared for drug from the standard stock solution and the solutions were scanned in the wavelength range of 400-800 nm. Absorption maxima were found to be 561nm, which was selected as wavelength of analytical measurement for this method. Calibration curve Appropriate volume of aliquots from standard Efavirenz stock solution II were transferred to different volumetric flasks of 10 ml capacity. To all the volumetric flasks 1ml of 1% sodium nitrite, 1ml of HCl and finally *hhw* was added. The volume was adjusted to the mark with ethanol to obtain concentrations of 10, 12, 14, 16, 18 and 20. Absorbance of each solution blank was measured at 561 nm and the graph of absorbance against concentration was plotted and is shown in Figure 10. The regression equation and correlation coefficient were determined. The method was validated according to International Conference on Harmonization guidelines for validation of analytical procedures (Raman, et al. [9,10,11]). Efavirenz has the absorbance maxima at 561 nm for the first order derivative visible-Spectrophotometric method. The optical characteristics such as Beer's limits, Limit of detection and Limit of

quantification etc., in each method were calculated and the results were presented in Table 1 respectively. The regression characteristics like slope (b), intercept (a), and correlation coefficient (R2) using the method of least squares were calculated and presented in Table 1 respectively. The results showed that the methods have reasonable precise. The validation parameters like Precision and Accuracy results were presented in Table 2 and 3 for Visible Spectrophotometric method. There was no interference of excipients in recovery study of the method. From results, the method found applicable for both bulk and pharmaceutical dosage for the estimation of Efavirenz. Recovery studies The recovery studies were carried out at three different level i.e. 80,100 and 120%. It was performed by adding known amount of standard drug solutions of Efavirenz to preanalysed tablets solutions. The resulting solutions were then reanalyzed by proposed methods. The results of recovery studies are shown in Table 4 and 5.




Table No.1 - Optical characteristic

1	Detection wavelength (nm)	561
2	Beer's law limits (µg/ml)	10-20
3	Regression equation (y*)	$y = 0.038x - 0.001$
4	Slope (b)	0.038
5	Intercept (a)	0.001
6	Correlation coefficient (R ²)	0.999
7	Limit of Detection (µg/ml)	0.078
8	Limit of Quantification	0.236

Table No.2 - Determination of Accuracy results for Efavirenz

Amount of sample (µg/ml)	Amount of drug added (µg/ml)	Amount Recovered(µg/ml)	% Recovery ± SD
16	12	11.98	99.78±0.46
16	16	15.89	99.08±0.95
16	20	19.99	99.97±0.11

Table No.4- Results of recovery studies for Efavirenz

Amount Of Sample($\mu\text{g/ml}$)	Amount of drug added ($\mu\text{g/ml}$)	% Recovery \pm RSD	%RSD
16	8	99.10 \pm 0.41	0.41
16	10	99.58 \pm 0.60	0.64
16	12	98.80 \pm 0.88	0.89

Table No. 5- Results of recovery studies for Sustiva

Amount Of Sample($\mu\text{g/ml}$)	Amount of drug added($\mu\text{g/ml}$)	% Recovery \pm RSD	%RSD
16	8	98.88 \pm 0.70	0.74
16	10	99.75 \pm 0.74	0.78
16	12	98.80 \pm 0.51	0.56

References

- Ribeiro JA, de Campos LM, Alves RJ, Lages GP, Pianetti GA (2007) Efavirenz related compounds preparation by hydrolysis procedure: Setting reference standards for chromatographic purity analysis. *J Pharm Biomed Anal* 43: 298-303.
- Montgomery ER, Edmanson AL, Cook SC, Hovsepiyan PK (2001) Development and validation of a reverse-phase HPLC method for analysis of efavirenz and its related substances in the drug substance and in a capsule formulation. *J Pharm Biomed Anal* 25: 267-284.
- Weissburg RP, Montgomery ER, Junnier LA, Segretario J, Cook S, et al. (2002) Investigation of critical factors for the resolution of SR695, a key impurity, from efavirenz in the reversed-phase assay of efavirenz dosage forms. *J Pharm Biomed Anal* 28: 45-56.
- Maurin MB, Rowe SM, Blom K, Pierce ME (2002) Kinetics and mechanism of hydrolysis of efavirenz. *Pharm Res* 19: 517-521.
- Gadkari T, Chandrachud P, Shahajetele A, Deshpande N, Salvekar J, et al. (2010) Validated stability indicating LC-PDAMS method to investigate pH rate profile and degradation kinetics of efavirenz and identification of hydrolysis products by LC-MS. *Int J Pharm Sci* 2: 169-176.
- Efavirenz monograph (2015) United States Pharmacopeia USP38, NF33, United States Pharmacopoeial Convention Inc: Rockville, CA, USA, p. 3656.
- Efavirenz tablets monograph (2015) United States Pharmacopeia, USP38, NF33, United States Pharmacopoeial Convention Inc, Rockville, CA, USA, p. 3661.
- Efavirenz capsules monograph (2015) United States Pharmacopeia, USP38, NF33, United States Pharmacopoeial Convention Inc, Rockville, CA, USA, p. 3659.
- Raman NVVSS, Prasad AVSS, Reddy KR (2011) Strategies for the identification, control and determination of genotoxic impurities in drug substances: A pharmaceutical industry perspective. *J Pharm Biomed Anal* 55: 662-667.
- Dow LK, Hansen MM, Pack BW, Page TJ, Baertschi SW (2013) The assessment of impurities for genotoxic potential and subsequent control in drug substance and drug

product. *J Pharm Sci* 102: 1404-1418.

- Robinson DI (2010) Control of genotoxic impurities in active pharmaceutical ingredients: A review and perspective. *Org Proc Res Dev* 14: 946-959.
- Elder DP, Lipczynski AM, Teasdale A (2008) Control and analysis of alkyl and benzyl halides and other related reactive organohalides as potential genotoxic impurities in active pharmaceutical ingredients (APIs). *J Pharm Biomed Anal* 48: 497-507.
- Bolta HM, Fothb H, Hengstlerc JG, Degena GH (2004) Carcinogenicity categorization of chemicals-New aspects to be considered in a European perspective. *Toxicol Lett* 151: 29-41.
- Boyle GA, Edlin CD, Li YF, Morgans GL, Musonda CC (2012) Enantioselective synthesis of the carbocyclic nucleoside abacavir [J]. *Org Biomol Chem* 10: 1870-1876.
- Shibata T, Morioka H, Hayase T, Choji K, Soaie K (1996) Highly enantioselective catalytic asymmetric automultiplication of chiral pyrimidyl alcohol [J]. *J Am Chem Soc* 118: 471-472
- Tan L, Chen CY, Tillyer RD, Grabowski EJJ, Reider PJ (1999) A novel, highly enantioselective ketone alkynylation reaction mediated by chiral zinc aminoalkoxides [J]. *Angew Chem Int Ed* 38(5): 711-713
- Pierce ME, Parsons RL, Radesca LA, Young SLO, Silverman S, et al. (1998) Practical asymmetric synthesis of efavirenz (DMP 266), an HIV-1 reverse transcriptase inhibitor [J]. *J Org Chem* 63: 8536-8543.

Assets of Publishing with us

Global archiving of articles
 Immediate, unrestricted
 online access Rigorous Peer
 Review Process Authors
 Retain Copyrights

<https://www.biomedress.com>

Submission Link: <https://biomedress.com/online-submission.php>