

Contribution of Immunoglobulin receptor (FcγR) and proteins of complement system in neutrophil to pathogenesis and progression of Rheumatoid Arthritis

Introduction

RA patients' blood neutrophils are functionally very different from healthy individuals' blood neutrophils: the former are primed for ROS production whilst the latter exist in a resting state in the circulation. These types of neutrophils also differ strikingly in their gene and protein expression patterns [1]. Systemic activation of the alternative complement pathway in active RA patients affects the effector functions of peripheral blood neutrophils [2]. Neutrophils from active RA and inactive RA patients responded to an exogenous complement source with significantly increased ROS production; however, incubating RA neutrophils with autologous sera indicated that there was poor FcγR/CR cooperation [2]. These results suggest that differences in the oxidative metabolism of neutrophils may reflect an acquired characteristic of the disease associated with distinct clinical manifestations.

- a) It is well established that simultaneous binding of opsonized particles to complement and Fcγ receptors enhances the efficiency of particle phagocytosis by immune cells including neutrophils. CR1 and CD32 expressed on neutrophils promote IC adherence, thereby increasing phagocytosis, oxidative burst and neutrophil degranulation compared to IC interaction with each receptor alone, providing evidence of synergy between these receptors [2].
- b) In a study of RA patients stratified according to disease activity, the level of FcγRIIIa and CR1 expression in active RA patients was higher compared to healthy controls and inactive RA patients. The cooperative ligation of these receptors led to increased O₂⁻ production by neutrophils from patients with active RA. This is important because ROS are harmful mediators of autoimmune inflammation in the joints [2].
- c) More recently, it was shown that the cooperation between C5aR and FcγRs is very important to initiate and sustain neutrophil recruitment in vivo in an animal model. Specifically, C5aR activation of neutrophils is required for LTB₄ release and early neutrophil recruitment to the joint, whereas FcγR engagement of neutrophils induces IL-1β release, ensuring continued inflammation [3]. These data

Open Access

Short Communication

Larissa Fávaro Marchi*

Centro Universitário Padre Albino-UNIFIPA, Catanduva, Brazil

*Address for Correspondence

Larissa Fávaro Marchi, Centro Universitário Padre Albino-UNIFIPA, Catanduva, Brazil

Submission: July 18, 2019

Published: July 24, 2019

Copyright: ©  This work is licensed under Creative Commons Attribution 4.0 License

support the concept that IC-mediated leukocyte activation is not composed of overlapping and redundant pathways but that each element serves a distinct and critical function in vivo, culminating in tissue inflammation. However, studies have highlighted the importance of CR in inflammation of articulated sites. Mice lacking the C5aR or C3aR have milder disease in arthritis models, showing that at least a part of the arthritis pathology is related to anaphylatoxin production and the interaction between anaphylatoxins and their cellular receptors [4,5].

- d) Our research team has contributed to understand the mechanism of action of antioxidant compounds on neutrophils and to develop therapeutic molecules that can modulate neutrophil ROS production in inflammatory diseases and be used in the complementary therapy in RA patients. We have investigated the in vitro pharmacological effects of natural and synthetic compounds such as flavonoids [6-8], coumarins [9,10] and sesquiterpene lactones [11] on the neutrophil ROS production and lysosomal enzyme release mediated by FcγR and CR.
- f) In the last decade, a lot of mechanisms of molecular and cellular signal transduction involved in IC deposition and in the inflammatory process have been unraveled. This information has contributed to the strategic development of modern therapies to treat inflammatory diseases in which the neutrophil has emerged as an important target of novel drugs.

References

1. Marzocchi-Machado CM, Alves CMOS, Azzolini AECS, Polizello ACM, Carvalho IF, et al. (2002) Fc gamma and complement receptors: expression, role and co-operation in mediating the oxidative burst and degranulation of neutrophils of Brazilian systemic lupus erythematosus patients. *Lupus* 11: 240-248.
2. Paoliello-Paschoalato AB, Moreira MR, Azzolini AECS,

- Cavenaghi AA, Marzocchi-Machado CM, et al. (2011) Activation of Complement Alternative Pathway in Rheumatoid Arthritis: Implications in Peripheral Neutrophils Functions. *Open Autoimmun J* 3: 1-9.
3. Alves CMOS, Marzocchi-Machado CM, Louzada-Junior P, Azzolini AECS, Polizello ACM, et al. (2008) Superoxide anion production by neutrophils is associated with prevalent clinical manifestations in systemic lupus erythematosus. *Clin Rheumatol* 27: 701-708.
 4. Sadik CD, Kim ND, Iwakura Y, Luster AD (2012) Neutrophils orchestrate their own recruitment in murine arthritis through C5aR and Fc gamma R signaling. *Proc Natl Acad Sci USA* 109: E3177-E3185.
 5. Banda NK, Hyatt S, Antonioli AH, White JT, Glogowska M, et al. (2012) Role of C3a Receptors, C5a Receptors, and Complement Protein C6 Deficiency in Collagen Antibody-Induced Arthritis in Mice. *J Immunol* 188: 1469-1478.
 6. Kanashiro A, Kabeya LM, Martinello F, Turato WM, Paula FS, et al. (2007) Effect of rutin on polymorphonuclear leukocytes oxidative metabolism in hypercholesterolemic Golden Syrian hamsters: evaluation by chemiluminescence and flow cytometry. *Pharmazie* 62: 295-298.
 7. Santos EO, Kabeya LM, Figueiredo-Rinhel AS, Marchi LF, Andrade MF, et al. (2014) Flavonols modulate the effector functions of healthy individuals' immune complex-stimulated neutrophils: A therapeutic perspective for rheumatoid arthritis. *Int Immunopharmacol*: 102-111.
 8. Marchi LF, Paoliello-Paschoalato AB, Oliveira RDR, Azzolini AECS, Kabeya LM, et al. (2018) Activation status of peripheral blood neutrophils and the complement system in adult rheumatoid arthritis patients undergoing combined therapy with infliximab and methotrexate. *Rheumatol Int* 38:1043-1052.
 9. Kabeya LM, de Marchi AA, Kanashiro A, Lopes NP, da Silva CH, et al. (2007) Inhibition of horseradish peroxidase catalytic activity by new 3-phenylcoumarin derivatives: Synthesis and structure-activity relationships. *Bioorg Med Chem* 15: 1516-1524.
 10. Andrade MF, Kabeya LM, Azzolini AE, Santos EO, Figueiredo-Rinhel AS, et al. (2013) 3-Phenylcoumarin derivatives selectively modulate different steps of reactive oxygen species production by immune complex-stimulated human neutrophils. *Int Immunopharmacol* 15: 387-94.
 11. Kanashiro A, Kabeya LM, Graef CF, Jordão CO, Azzolini AE, et al. (2006) Sesquiterpene lactones from *Lychnophora pohillii*: Neutrophil chemiluminescence inhibition and free radical scavenger activity. *J Pharm Pharmacol* 58: 853-858.

Assets of Publishing with us

Global archiving of articles
Immediate, unrestricted
online access Rigorous Peer
Review Process Authors
Retain Copyrights

<https://www.biomedress.com>

Submission Link: <https://biomedress.com/online-submission.php>